



# Studies of Physiological and Yield Attributing Traits of Soybean [*Glycine max* (L.) Merrill]

Akash Barela <sup>a\*</sup>, M. K. Shrivastava <sup>a</sup>, Yogendra Singh <sup>a</sup>,  
Pawan K. Amrate <sup>a</sup>, Salikram Mohare <sup>b</sup>, Shivani Jawarkar <sup>a</sup>  
and Satish Kumar Nagar <sup>a</sup>

<sup>a</sup> Department of Plant Breeding and Genetics, JNKVV, Jabalpur 482004, M.P., India.

<sup>b</sup> Department of Plant Physiology, JNKVV, Jabalpur 482004, M.P., India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i193552

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104585>

Original Research Article

Received: 06/06/2023

Accepted: 10/08/2023

Published: 18/08/2023

## ABSTRACT

The experiment was conducted at AICRP Soybean, Seed Breeding Farm, Department of Plant Breeding and Genetics, College of Agriculture, JNKVV, Jabalpur (M.P.). The experiment was consisting of 90 genotypes to study the difference in growth, morpho-physiological characterization and yield response of soybean. The genotype JS 21-72 followed by JS 22-18 and PS-1613 recorded the highest number of leaves per plant. AGS-48 showed maximum plant height among all other genotypes. Genotype DS 31-09 followed by JS 22-16 produced the maximum number of pods per plant. DS 31-09 followed by JS 22-16 produced maximum number of pods per plant with respects to all another genotype Early work with infrared thermometers proved successful in monitoring plant Evaporative rate. The canopy temperature in the current study ranged from 32.13

\*Corresponding author: E-mail: akashbarelagpb@gmail.com;

in JS 23-03 to 36.15 in JS 22-49. In yield attributing characters genotype highest seed yield was produced by JS 21-72 whereas JS 24-35 followed by JS 24-37 the shows highest number of seed per pod and 100 seed weight. Genotype NRC-137 followed by JSM 259 showed a higher range of harvest index. It is concluded that the six genotypes viz., JS 21-72, JS 24-35, JS 24-37, JS 22-16 AGS-48 and NRC- 137 found best to describe the morpho-physiological characterization in the present study.

**Keywords:** Canopy temperature; LAI; LAD and morpho-physiology.

## 1. INTRODUCTION

Soybean [*Glycine max* (L.) Merrill.] is one of the miraculous crops due to its extraordinary qualities. It has transformed the agricultural economy through its enormous potential for food, fuel, and a variety of industrial products. Its production has a very complex character; it is reliant on a variety of physiological and genetic elements that interact with the environment. Genetic yield potential is static as a result of unclear productivity restrictions, such as decreasing sink demand, which invites a huge gap in critical yield status. It is vitally necessary to determine the potential key physiological variables that are the physiological determinant and connected with the seed yield of crop in light of the rapid depletion of agricultural resources. This may be the sole glimmer of hope for choosing the genotype that is best for Madhya Pradesh. Currently, soybean has emerged as a key agricultural product, contributing to roughly 25% of the world's production of edible oil as well as two-thirds of the protein concentrate used in animal, poultry, and fish feed Agrawal et al., [1]. In India, soybeans are the main oilseed crop farmed in the Kharif season (10.93 mt production and 10.83 M Ha area), and the nation ranks fifth globally in soybean production behind the United States, Brazil, China, Argentina, and USA Anonymous [2]. In India, soybean genotypes and cultivars exhibit limited genetic diversity Agrawal et al., [1], Barela et al., [3] and Mehra et al., [4]. There are limitations of knowledge regarding the physiological growth characteristics (such as relative growth rate, leaf area ratio, specific leaf area ratio, and shoot/root) that correspond to the responses of various soybean genotypes to various plant densities.

## 2. MATERIALS AND METHODS

The present investigation was carried out during kharif 2021, it was conducted at AICRP Soybean, Seed Breeding Farm, Department of Plant Breeding and Genetics, College of Agriculture, JNKVV, Jabalpur (M.P.). The

experiment was laid out in Randomized Block Design (RBD) with three replications and 90 genotypes. The experimental site had uniform topography, medium black soil with 7.5 pH and was free from water-logged conditions. Leaf area index is the leaf area existing on unit ground was proposed by Watson [5] and was calculated by the following formula:

$$LAI = \frac{\text{Leaf area per plant (cm}^2\text{)}}{\text{Ground area per plant (cm}^2\text{)}}$$

Power et al., (1967) integrated LAI with time and this was known as leaf area duration. LAD takes into account both the duration and the extent of photosynthetic tissue of the crop canopy. The LAD is expressed in days. The LAD was calculated for 60 DAS and 90 DAS as LAD1 and LAD 2 respectively using the formula as follows:

$$LAD = \frac{(LA2+LA1) \times (t2-t1)}{2}$$

Where

LA1 and LA2 represent the leaf area at two successive time intervals (t1 and t2)

Harvest index (HI) was determined from the mean value of seed yield per hectare and biological yield per hectare at the time of harvest using a formula given by Donald [6].


$$\text{Harvest Index} = \frac{\text{Economic yield}}{\text{Biological yield}} \times 100$$

## 3. RESULTS AND DISCUSSION

The results highlighted the increased trends in plant height, which is statistically significant at all the growth stages of the crop (Table 1). JS 22-41 followed by JS 22-43 and JSM-259 recorded a significant maximum plant height at 60 DAS (Days After Sowing) whereas AGS-48 followed by JS 22-04 and JS 22-52 recorded the maximum height at 90 DAS. In the current investigation, it was evident that the genotype with short stature showed more tolerance against

drought as compared to the taller once Mishra et al., [7] Genotypic differences in plant height in soybean were also reported by Abbas et al. [8]. Genotype JS 21-72 followed by DS 31-09, JS 22-16 produced the maximum number of pods per plant with respect to all other genotypes and these results are in conformity with Deshmukh et al. [9] in soybean. Genotype JS 21-72 followed by JS 22-18 and PS-1613 recorded the highest number of leaves per plant (Table 1). The differences in mean leaf area were statistically significant at 60 and 90 DAS (Table 1). The genotype JS 24-37 followed by JS 22-26 and JS 22-33 maintained the highest value of leaf area from 60 DAS onward. Similar result was observed by Gontia and Awasthi [10] in soybean. LAI and biomass are the two key indicators of plant health and development Barela et al., [3]. The growth analytical parameter studies like Canopy temperature, LAI and LAD exhibited significant trends (Table 1). According to recent studies infrared thermometers have been a useful instrument for measuring canopy temperatures in semi-arid and dry settings. The optimal canopy temperatures for peak photosynthetic enzyme activity, which were discovered to be 27°C, served as the basis for the soybean temperature thresholds (Fig. 2). Early work with infrared thermometers proved successful in monitoring plant Evaporative rate, the canopy temperature in the current study ranged from 32.13 in JS 23-03 to 36.15 in JS 22-49. The genotypic difference with respect to

Canopy temperature, LAI and LAD were reported by Rahangdale et al., [11], Rajput and Shrivastava [12] in soybean. All the genotypes were different in yield attributing character (Table 1). As regards to yield attributing Table 1: Morpho-physiological characters of soybean genotype included Plant height at 30,60 and 90 DAS (Days after sowing), leaf number at 30,60 and 90 DAS, LAI(Leaf area index) at 60 and 90 DAS, LAD (Leaf area duration at 60 and 90 DAS), CT(Canopy temperature), S/POD(Seed per pod), P/P (Pods per plant), NS/P(Number of seeds per plant), 100SW(Hundred seeds weight), BY/P(Biological yield per plant), HI(Harvest index),SY/P(Seed yield per plant). Similar result was obtained by Singh et al. [13]. The genotype JS 24-35 followed by JS 24-37 recorded the highest number of seeds per pods and 100 seed weight. Genetic variability in soybean was also obtained by Singh et al. [13]. The genotype JS 21-72 followed by JS 22-16 recorded highest seed yield per plant and biological yield per plant as well. Seed per pod were highest in JS 22-52 followed by JS 22-32 and JS 22-33, whereas the highest seed yield produced by JS 21-72 might be due to highest number of pods per plant and number of seed per plant. Similar variations in soybean were also obtained by Deshmukh et al. [9]. The genotype NRC-137 followed by JSM 259 showed highest value of harvest index and this in conformity with the results of Nirmal Kumari and Balasubramanian [14] in soybean.



**Fig. 1. Graphical representation of Leaf Area Index in different series of LAI at 60 DAS and 90 DAS respectively**

**Table 1. General mean values of each genotype for physiological and quantitative traits**

Germplasm	Plant height			Leaf number			Leaf area index		Leaf area duration			S/ POD	P/P	NS/P	100 SW	BY/P	HI	SY/P
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	60 DAS	90 DAS	LAD1	LAD2	CT							
1 AGS 48	17.10	32.92	52.17	21.33	66.33	79.33	6.56	9.08	66.06	83.68	33.18	2.17	38.00	116.50	9.32	25.29	47.57	12.09
2 AGS 163	20.67	28.45	40.33	22.67	61.67	86.00	6.07	7.70	73.28	81.48	34.17	2.33	46.03	121.67	10.73	22.43	60.18	13.49
3 AVKS 199	19.00	26.19	38.00	22.67	65.33	86.00	6.41	7.92	67.14	84.67	34.10	2.17	39.50	121.83	11.58	22.84	58.52	13.32
4 AVKS 206	19.53	25.77	41.33	22.33	67.00	80.33	7.63	8.89	70.49	82.19	35.35	2.33	41.00	68.83	10.90	14.23	56.17	8.09
5 Cat 87	20.02	25.30	36.50	23.33	65.00	69.67	5.59	6.82	74.50	85.87	34.47	2.50	37.00	68.50	10.53	16.34	45.99	7.46
6 Cat 783	19.19	24.27	32.67	24.00	64.67	80.33	7.56	7.92	70.49	81.47	33.40	2.00	38.50	100.50	11.07	14.78	74.70	10.72
7 JS 24-35	17.53	26.97	33.33	25.00	63.67	82.33	6.53	8.89	67.93	82.11	33.90	3.00	34.67	133.67	10.35	20.74	58.37	12.10
8 DS 3106	19.20	29.21	32.83	24.67	65.00	80.00	6.26	8.98	69.71	83.15	33.03	2.33	28.50	61.00	10.10	14.17	44.75	6.32
9 DS 3109	19.03	24.33	39.00	22.00	61.33	76.00	5.38	7.13	73.70	92.39	33.10	2.50	51.00	86.67	11.35	16.57	57.97	9.62
10 JS 24-36	17.70	26.19	42.17	24.33	63.67	76.00	5.82	8.19	68.85	79.00	33.19	2.50	32.50	54.17	12.10	16.47	49.45	8.12
11 EC350664	19.17	29.01	35.00	20.33	64.33	78.33	6.85	7.20	69.45	87.22	33.93	2.17	37.67	68.17	10.78	18.17	44.35	7.99
12 EC481571	19.77	24.33	38.17	21.33	63.33	79.33	6.41	8.24	66.13	90.26	33.57	2.67	36.17	84.00	10.83	24.28	40.47	9.84
13 JS 335	18.60	26.19	34.50	25.67	62.00	78.33	6.85	10.00	69.03	82.07	33.55	2.67	32.50	51.83	11.10	26.06	30.90	7.75
14 JS 2069	18.06	25.33	31.00	23.67	62.67	87.67	6.41	9.22	67.71	80.70	33.64	2.67	34.17	82.83	10.20	24.59	37.02	9.14
15 JS 20-73	19.87	29.21	33.33	22.67	62.33	81.67	6.68	6.82	70.35	86.55	34.34	2.67	36.00	89.50	10.70	21.52	44.54	9.57
16 JS 20-98	19.37	24.33	30.00	24.33	65.00	75.67	7.56	7.92	74.71	90.18	34.08	2.17	38.67	106.50	9.40	16.74	58.75	9.81
17 JS 20-105	18.58	26.72	33.50	23.00	64.00	78.00	6.53	8.57	69.58	89.91	35.08	2.67	47.44	119.50	12.05	26.12	51.77	13.28
18 JS 20-116	17.71	27.22	39.17	20.67	62.00	77.67	6.26	8.98	66.06	89.09	34.17	2.44	45.81	106.17	9.10	22.30	52.28	11.62
19 JS 21-05	19.82	26.19	38.83	23.33	59.33	77.33	5.30	8.94	73.28	85.73	34.10	2.83	31.67	70.67	11.61	16.32	55.73	9.28
20 JS 21-72	19.35	26.19	38.33	25.67	68.00	89.00	7.71	9.44	71.75	76.73	34.73	2.33	52.57	132.00	12.63	25.00	69.38	17.35
21 JS 22-03	20.15	26.19	41.17	23.00	64.33	74.33	6.05	7.12	64.53	88.72	33.24	2.50	29.62	86.17	10.80	20.74	46.29	9.59
22 JS 22-04	19.08	24.67	51.83	23.67	65.33	74.67	5.80	7.92	67.97	90.53	35.18	2.33	25.40	54.00	11.55	14.17	50.59	7.17
23 JS 22-10	18.38	26.05	33.67	24.33	59.67	75.67	6.94	8.98	67.21	84.77	34.27	2.17	30.67	68.83	9.85	16.57	42.98	7.15
24 JS 22-11	17.91	25.70	36.00	23.67	64.00	89.00	6.42	8.61	67.46	84.22	34.28	2.83	24.34	67.67	12.80	21.52	36.77	7.91
25 JS 22-12	20.67	24.33	36.33	21.33	67.67	82.67	6.24	7.94	69.03	90.53	34.00	2.33	31.26	107.33	10.86	20.74	61.90	12.87
26 JS 22-13	18.53	29.61	36.83	23.33	64.67	87.33	6.62	8.25	77.38	85.33	35.08	2.83	28.28	82.00	10.17	14.17	63.93	9.05
27 JS 22-14	17.53	24.67	38.67	23.00	65.67	82.67	5.66	7.92	66.06	87.78	34.10	3.00	24.45	72.50	8.85	16.06	41.54	6.73
28 JS 22-15	18.57	24.74	44.33	24.67	63.33	79.33	6.85	8.57	73.28	92.39	35.35	2.83	43.70	110.17	13.10	22.90	56.42	12.91
29 JS 22-16	19.30	24.30	40.50	23.00	66.33	83.00	6.41	8.98	77.19	84.06	34.47	2.50	50.47	127.89	11.73	24.90	63.48	15.22
30 JS 22-17	17.10	26.60	38.33	23.33	64.33	80.33	7.63	8.94	78.13	87.15	33.40	2.33	34.67	62.83	12.44	20.74	47.64	9.85
31 JS 22-18	20.67	25.03	36.00	25.33	66.70	88.33	4.58	7.88	74.70	92.39	33.90	3.00	25.83	59.00	10.61	20.74	38.81	8.04
32 JS 22-19	17.00	24.74	37.00	26.33	64.00	75.67	5.96	7.88	70.35	90.86	33.03	2.67	27.45	83.00	10.48	17.24	47.49	8.14
33 JS 22-20	19.20	24.30	46.67	22.00	65.67	85.67	5.32	7.85	74.71	81.48	33.10	2.50	31.84	50.67	9.50	16.57	44.81	7.41
34 JS 22-22	18.03	26.60	40.67	22.33	66.33	81.00	7.61	7.95	75.84	90.18	33.19	3.00	32.50	67.00	11.80	20.09	38.36	7.45
35 JS 22-23	20.20	28.38	35.17	25.33	63.00	84.67	6.24	8.94	69.30	80.70	33.93	3.00	28.67	62.17	13.00	21.52	34.05	7.32
36 JS 22-24	17.70	26.19	38.67	24.67	67.67	75.67	7.51	7.88	69.03	85.61	36.15	2.67	32.83	41.17	11.45	22.84	40.48	9.22
37 JS 22-25	19.17	27.16	42.50	22.67	66.33	77.00	5.03	7.47	68.24	91.45	34.10	2.50	31.00	65.50	9.40	14.17	59.52	8.44
38 JS 22-26	17.53	26.19	35.50	24.00	67.00	78.00	7.63	10.10	63.74	88.04	35.08	2.50	28.50	55.22	11.60	21.52	35.63	7.67
39 JS 22-27	20.20	27.16	41.00	24.33	63.33	81.67	7.36	9.78	70.49	85.33	34.00	2.67	36.45	121.00	12.30	20.74	55.77	11.52
40 JS 22-28	19.00	23.37	33.50	23.67	64.33	79.67	6.24	7.95	74.31	94.57	35.08	2.50	20.67	59.33	11.45	22.43	30.51	6.84

	Germplasm	Plant height			Leaf number			Leaf area index		Leaf area duration			S/ POD	P/P	NS/P	100 SW	BY/P	HI	SY/P
		30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	60 DAS	90 DAS	LAD1	LAD2	CT							
41	JS 22-29	19.53	26.19	34.33	22.00	63.67	83.67	7.22	8.94	73.45	88.97	34.55	2.33	19.50	53.00	10.10	20.74	26.73	5.55
42	JS 22-30	20.02	28.35	37.00	24.00	65.33	82.00	6.24	7.88	67.97	78.58	35.17	2.83	38.00	111.67	11.33	16.37	92.14	12.71
43	JS 22-31	19.19	26.19	39.67	23.67	64.33	85.67	6.62	7.47	73.56	85.61	33.98	2.83	26.67	51.17	11.70	24.59	25.88	6.34
44	JS 22-32	17.53	27.35	35.17	19.00	67.67	80.00	5.66	8.23	73.60	91.45	33.03	3.10	30.33	59.33	12.15	21.52	38.64	8.32
45	JS 22-33	19.20	29.01	46.67	24.33	66.67	71.00	7.95	10.08	66.59	90.86	32.67	3.07	22.83	51.33	11.69	20.74	30.29	6.28
46	JS 22-34	19.50	24.67	39.50	24.33	63.67	83.33	6.82	9.22	70.35	85.63	33.86	2.67	32.83	87.00	10.63	18.74	49.39	9.70
47	JS 22-35	20.20	27.22	37.83	20.67	69.33	78.67	6.53	6.46	74.71	85.61	33.77	2.83	37.33	55.00	10.55	16.57	46.71	7.71
48	JS 22-36	17.10	26.19	38.50	22.33	65.00	84.33	7.63	9.58	75.84	85.93	34.60	2.83	37.84	101.17	9.65	32.65	33.88	11.06
49	JS 22-38	20.67	26.19	35.00	20.33	60.33	84.67	5.59	8.98	70.20	83.67	34.16	2.17	30.50	60.00	12.65	20.18	38.70	7.52
50	JS 22-39	20.33	26.19	41.67	21.33	64.33	83.67	6.62	8.50	67.46	78.18	33.42	2.83	35.00	118.17	8.60	16.32	68.54	10.70
51	JS 22-40	19.91	24.67	45.50	26.00	65.33	78.67	5.63	9.68	69.03	80.70	34.85	2.50	32.50	62.00	10.60	21.52	30.41	6.55
52	JS 22-41	18.80	29.87	35.83	25.00	64.33	74.33	5.90	8.15	77.38	84.97	33.42	2.50	26.50	56.00	10.40	20.74	33.76	6.99
53	JS 22-42	17.23	24.33	29.33	22.33	63.67	86.00	7.51	8.24	70.49	89.91	35.67	2.83	32.67	50.50	12.80	29.65	29.58	8.37
54	JS 22-43	18.70	29.61	36.00	25.00	63.67	80.67	6.79	8.98	67.93	90.53	33.42	2.50	24.67	62.67	12.50	21.52	37.72	8.12
55	JS 22-44	19.40	24.67	35.67	20.33	66.00	91.00	5.96	8.26	76.12	84.97	35.17	2.50	25.67	64.33	11.65	22.43	35.31	7.92
56	JS 22-45	17.00	24.74	49.67	21.67	60.33	83.67	5.92	7.85	73.60	96.38	35.08	2.83	32.00	51.50	11.20	23.89	34.23	7.46
57	JS 22-46	18.40	24.74	31.67	21.00	65.67	78.33	5.54	7.92	63.74	85.93	34.10	2.50	25.50	55.50	10.10	16.57	45.64	7.59
58	JS 22-47	18.37	24.30	41.50	21.67	62.00	81.00	6.80	9.74	70.21	83.67	35.35	2.50	29.83	56.33	11.25	21.52	32.59	7.02
59	JS 22-48	16.90	26.60	51.00	23.67	67.33	77.00	6.85	10.00	70.43	89.91	34.47	2.83	29.17	72.33	11.00	21.52	39.57	8.50
60	JS 22-49	20.67	25.03	33.67	22.00	60.00	85.00	5.54	9.22	67.71	90.53	36.15	2.17	45.00	103.50	12.80	23.67	54.56	13.03
61	JS 22-50	17.00	24.74	32.33	20.67	64.00	72.67	7.51	7.21	73.28	86.79	34.10	2.50	35.50	66.33	10.60	14.17	64.60	9.14
62	JS 22-51	19.20	24.30	33.00	24.67	63.67	88.00	6.79	8.98	69.35	89.97	34.34	2.83	29.00	60.00	10.34	16.57	43.61	7.19
63	JS 22-52	18.20	26.60	51.67	20.33	66.00	80.67	7.19	8.50	75.84	83.63	33.10	3.17	31.17	67.83	12.70	20.74	41.07	8.52
64	JS 22-53	19.67	28.38	33.33	25.00	63.00	76.00	5.88	9.68	70.20	84.97	33.19	2.50	37.33	70.67	12.60	20.18	45.23	9.13
65	JS 22-54	19.87	26.19	42.67	23.33	66.00	84.33	6.85	8.98	75.93	90.86	33.93	2.50	33.83	67.00	13.10	23.40	38.53	9.03
66	JS 22-55	16.90	27.35	37.17	24.67	62.33	84.67	6.29	7.95	67.97	81.57	33.57	3.00	26.00	59.67	11.85	21.52	32.37	6.96
67	JSM 259	18.20	29.01	40.50	24.67	60.33	75.33	5.88	8.24	67.55	84.97	33.40	2.83	41.50	89.83	12.65	21.74	67.21	14.14
68	JSM 232	19.20	24.67	33.00	22.33	64.33	78.33	5.69	8.98	75.10	88.31	33.97	2.83	32.67	90.00	12.55	16.16	67.03	10.75
69	JS 23-03	18.03	27.22	31.17	24.33	64.33	84.00	5.15	7.13	78.79	84.97	32.13	2.33	32.50	71.33	13.81	32.65	35.87	11.71
70	JS 23-05	17.23	27.16	39.17	24.33	64.67	86.00	6.85	8.88	71.67	88.04	33.40	2.33	29.67	78.00	13.02	29.65	36.59	10.74
71	JS 23-08	17.43	26.19	41.33	25.33	64.67	82.00	5.88	6.89	67.80	90.79	34.88	3.00	28.71	55.67	12.20	21.52	45.48	9.80
72	JS 23-09	18.53	26.19	30.33	22.00	65.00	81.67	6.85	9.58	73.21	84.97	33.26	2.17	33.00	87.00	12.40	22.43	53.27	11.95
73	JS 23-10	16.90	29.01	31.67	21.33	66.33	77.00	5.54	8.98	76.12	94.45	32.66	2.50	33.00	90.83	10.90	20.18	58.72	11.73
74	MAUS 71	17.70	26.19	33.33	23.67	61.00	82.00	7.51	7.13	74.82	92.39	34.83	2.67	34.33	86.33	13.47	16.57	49.52	8.25
75	MACS 450	20.67	24.67	32.67	24.00	68.67	82.00	7.57	9.68	67.93	84.97	34.27	2.50	28.17	50.83	10.90	21.52	43.40	9.34
76	NRC 125	18.70	27.42	33.67	22.67	68.33	80.67	5.73	7.88	76.12	88.04	33.42	3.00	44.17	100.83	13.25	22.67	59.46	13.56
77	NRC 128	19.67	29.01	37.00	21.33	65.67	81.67	6.24	10.00	74.82	84.97	34.22	2.67	32.00	67.00	11.95	14.17	55.10	7.79
78	NRC 137	17.10	24.67	34.50	24.00	65.67	78.67	5.48	8.98	73.21	92.39	34.00	2.83	42.00	120.00	12.52	18.57	68.03	13.56
79	PS 1092	20.67	24.40	31.00	24.67	65.67	84.33	5.61	8.58	75.10	84.97	35.08	2.50	31.50	53.67	12.45	16.72	44.00	7.36
80	PS 1225	19.91	25.70	33.50	25.00	65.67	89.33	7.88	7.95	78.79	92.39	34.27	2.67	31.67	64.83	12.73	29.65	31.05	8.98
81	PS 1613	18.80	29.21	33.50	26.30	68.33	86.67	5.22	8.94	76.12	84.97	34.00	2.67	31.00	67.33	12.05	21.52	43.07	9.27
82	PS 1641	17.23	24.33	30.33	23.33	64.33	82.33	6.91	7.88	67.97	88.04	35.08	2.67	37.56	105.17	11.44	18.00	62.29	11.24

Germplasm	Plant height			Leaf number			Leaf area index		Leaf area duration			S/POD	P/P	NS/P	100 SW	BY/P	HI	SY/P
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	60 DAS	90 DAS	LAD1	LAD2	CT							
<b>83 RVS 2012-19</b>	18.70	24.74	40.00	21.00	65.00	85.00	5.83	7.37	67.55	84.09	34.83	3.00	30.00	63.00	12.70	20.18	44.98	8.85
<b>84 SL 96</b>	19.50	25.42	31.50	21.33	62.00	82.67	7.30	9.37	75.10	84.97	34.27	2.67	33.50	56.50	10.34	16.32	49.46	7.70
<b>85 SL 710</b>	17.23	23.62	42.50	22.33	67.33	88.00	6.78	9.00	78.79	84.97	34.57	2.50	28.33	54.33	11.05	32.65	20.74	6.77
<b>86 SL 744</b>	17.20	27.30	37.17	23.67	66.67	89.00	5.92	8.98	76.12	92.39	35.08	2.67	28.33	58.67	10.73	20.74	35.40	7.34
<b>87 JS 24-37</b>	19.87	24.33	33.00	23.33	65.33	87.00	8.3.63	10.94	67.97	90.78	34.10	2.72	40.50	129.83	12.60	24.91	60.47	15.03
<b>88 NRC 86</b>	18.90	25.42	46.83	21.67	60.67	84.33	7.36	10.69	72.00	80.70	35.35	2.97	37.00	64.83	12.20	24.22	32.38	7.85
<b>89 NRC 130</b>	18.07	23.62	42.17	23.00	66.00	83.00	6.24	6.88	75.10	84.97	34.47	2.92	32.17	107.33	10.68	23.00	54.80	12.61
<b>90 PS 1569</b>	18.31	27.30	38.00	22.33	65.33	81.00	7.51	8.78	77.66	95.65	34.55	2.60	34.33	118.00	11.97	21.67	66.98	14.50
<b>Mean</b>	18.76	26.23	37.60	22.93	64.43	80.35	6.41	8.47	71.47	86.76	34.15	2.61	33.48	77.93	11.38	20.73	47.44	9.51
<b>SE</b>	0.83	1.76	1.90	1.58	2.00	4.16	0.96	0.88	3.71	5.18	0.61	0.15	2.08	6.86	0.31	1.33	5.51	0.85
<b>C.D.5%</b>	2.32		5.30						8.59		1.71	0.41	5.80	19.14	0.86	3.71	15.37	2.39

Note: Plant height at 30,60 and 90 DAS(Days after sowing), leaf number at 30,60 and 90 DAS, LAI(Leaf area index) at 60 and 90 DAS, LAD(Leaf area duration at 60 and 90 DAS), CT(Canopy temperature), S/POD (Seed per pod), P/P(Pods per plant), NS/P(Number of seeds per plant), 100SW(Hundred seeds weight), BY/P(Biological yield per plant), HI(Harvest index),SY/P(Seed yield per plant)




Fig. 2. Graphical representation of different physiological along with quantitative traits

#### 4. CONCLUSION

The observed results imply that physiological features restrict soybean genotypes' ability to develop and function metabolically. These features demonstrated the existence of significant diversity throughout several growth periods. The results of this work may help to identify various adaptive mechanisms developed by soybean genotypes and contribute to the identification of useful.

#### CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Agarwal DK, Billore SD, Sharma AN. Soybean: Introduction, improvement and utilization in India problems and prospects. *Agricultural Research*. 2013;2:293-3006.
2. Anonymous. The soybean processors association of India (SOPA) Report; 2020.
3. Barela A, Shrivastava MK, Mohare S, Rahangdale S, Jawarkar S, Amrate PK and Singh Y. Morphological Characterization and Recognition of New Traits of Soybean [*Glycine max* (L.) Merrill] *International Journal Environment and Climate Change*. 2022;12(12):1497-1504.
4. Mehra S, Shrivastava MK, Amrate PK, Yadav RB. Studies on variability, correlation coefficient and path analysis for yield associated traits in soybean [*Glycine max* (L.) Merrill]. *Journal of Oilseeds Research*. 2020;37(1):56–59.
5. Watson DJ. Physiological aspects of crop yield. *Adv. in Agron*. 1952;4:1001-1045.
6. Donald. In search of yield. *J. Australian agric. Sci*. 1962;28:171-178.
7. Fisher. Laboratory manual of physiological studies of rice. *IRRI*. 1921;30.
7. Mishra N, Tripathi MK, Tiwari S, Tripathi N, Trivedi HK. Morphological and molecular screening of soybean genotypes against

- yellow mosaic virus disease. Legume Research an International Journal; 2020. DOI: 10.18805/LR4240.
8. Abbas Mohd, Singh MP, Nigam KB, Kandalkar VS. Effect of plant densities and plant type on different growth and physiological parameters of soybean. Indian J. Agron. 1992;39(2):246-248.
  9. Deshmukh DT, Dhumale DR, Khordge PW, Kshirsagar AR. Inter- relationship and path analysis of yield and its components in soybean. Ann. Plant Physiol. 1991;6(1): 139-141.
  10. Gontia AS, Awasthi MK. Effect of seed size on various seed vigour attributes, morphological characters and seed yield in soybean genotype. Seed Res. 1998;27(1): 25-30.
  11. Rahangdale S, Lakhani JP, Singh SK, Barela A. Phenotyping and reveal novel traits in mungbean (*Vigna radiata* L. Wilczek) genotypes, South African Journal of Botany. 2023;156:65-71.
  12. Rajput RL, Shrivastava VK. Influence of varieties, sowing date and seed rate on physiological parameter and seed yield of soybean [*Glycine max* (L.) Merrill]. Legume Res. 1991;22(2):117-220.
  13. Singh SP, Rao SK, Shrama SM. Yield components in Soybean [*Glycine max* (L.) Merrill]. JNKVV Res. J. 1983;17(1-2):46-50.
  14. Nirmal Kumari A, Balasuramanian M. Variability in harvest index of soybean and its associated with yield components. Ann. Plant Physiol. 1991;5(2):256-258.

© 2023 Barela et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/104585>