



Monitoring Studies of Arthropod Complex Fauna Associated with Rapeseed- Mustard Crop Grown in Kashmir Himalayas

Muneer Ahmad Sofi ^{a*}, Bashir Rather ^a, Mushtaq Dar ^a,
Shafeeq hakeem ^b, Faisul Rasool ^c, Latief Ahmad ^c,
Aamir Hussain ^c and Bismat Un Nisa ^{d++}

^a MRCFC, Khudwani Kulgam, India.

^b Division of Entomology, Faculty of Horticulture, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, India.

^c Dry Land Agriculture Research Station, Rangreth, Srinagar, India.

^d Division of Entomology, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193579

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105091>

Original Research Article

Received: 15/06/2023

Accepted: 18/08/2023

Published: 22/08/2023

ABSTRACT

Kashmir has quietly undergone a yellow revolution as thousands of hectares of land have been brought under mustard cultivation to double the farmers' income. Earlier, most of the land in Kashmir was used only for growing a single crop -- paddy in most areas but now farmers are rotating crops in Kharif and Rabi seasons just like in other parts of the country. There is still many challenges which are hurdle in high production goal. Insect biodiversity is associated with the quantity and type of host plants available, environmental factors, and their physiological state.

**Ph.D Scholar;

*Corresponding author: E-mail: jabarimuneer@gmail.com;

Arthropod complex of mustard was studied and about eight species of insects-pests at Mountain Research centre for field crops (MRCFC) during Rabi 2020 -21 was recorded. It has been observed that the mustard aphid, *Lipaphis erysimi* Kalt., persists constantly in a sizable population and causes substantial crop damage from the blossoming to the maturity stage. *Phyllotreta cruciferae* Goeze (flea beetle) occurred at the seedling stage, and *Carpocoris* sp. (shield bug) caused little harm to the crop from the seedling to the maturity stage. Leaves have been only infested by cabbage aphid, *Brevicoryne brassicae*, mustard aphid, *Lipaphis erysimi* (Kalt.), and mustard leaf miner, *Chromatomyia horticola*, and they been detected in sporadic, low-population occurrences. Among them, *Brevicoryne brassicae* is found at flowering stage. *Pieris brassicae*, the cabbage butterfly, occurs sporadically and with high abundance. The status of all eight insect-pests on *Brassica* species was observed as minor with irregular occurrence while *L. erysimi* as a regular key pest. six Insect pollinators were recorded during the bloom period of the crop.

Keywords: Arthropod; rapeseed; mustard; insects; Kashmir; crop.

1. INTRODUCTION

“Amid a slew of initiatives by the government to double farmer’s income, mustard oilseed farming is gaining popularity in Kashmir as the farmers brought more land under its cultivation during 2020-21. The variety and availability of host plants, environmental variables, and physiological status are all related to insect biodiversity. As host plants serve as both acceptable oviposition sites for adult females and prospective food sources for the young ones, their availability and an insect's capacity to locate them are crucial for the survival of a species” [1]. “The information that the olfactory and gustatory systems transmit to the central nervous system (CNS) about the chemical composition of the plant plays a significant role in the choice and recognition of a host plant. Insects in particular exhibit a high degree of peripheral plasticity that enables them to adapt to their surroundings. In terms of production, rapeseed-mustard is the second most important oilseed crop farmed in India after soybean, and it has the highest oil yield among oilseed crops. It is produced on a 6.4 million hectare (ha) area with a productivity of 1262 kg/ha” [2]. “It has an oil content ranging from 35 to 45%. India produces 10% of the world's edible oils and holds the fourth-largest edible oil economy in the world, behind the US, China, and Brazil. The primary oilseed crop cultivated in India during rabi is rapeseed-mustard. The top states for mustard cultivation, accounting for more than 70% of the nation's total mustard area, are Rajasthan, Uttar Pradesh, Madhya Pradesh, Haryana, and Gujarat. *Brassica juncea*, one of the *Brassica* species, makes up roughly 90% of the total area in India. Even with the use of good production technologies, the nation's mustard crop is unable to produce its full potential output. This is owing

to the fact that different biotic and abiotic variables severely reduce the production of *Brassica* crops. Insect pests are one of the most significant biotic factors affecting crop productivity among biotic restrictions” [3]. In India, rapeseed-mustard has been discovered to be infested by about 50 different insect species. The Kashmir valley, where we have a moderate type of environment that is different from the rest of the country, offers significant potential for research on insect-pests of rapeseed and mustard [4-7].

2. MATERIALS AND METHODS

The study was conducted first time in Kashmir during rabi 2020-21 and 2021-22 in the Field and Laboratory of the Section of Entomology, Mountain research centre for field crops (MRCFC) Khudwani Kulgam SKUAST-K. The mustard *Brassica juncea*, variety Shalimar sarson -1 was collected from Breeding Centre and seeds were sown on 30 October in rows. The row to row distance was 45 cm. Intercultural operations such as weeding etc were done whenever necessary but no pest management practices were done in the field. Fertilizers were applied according as SKUAST-K Package and practices. Monitoring of the incidence of insect pests and their natural enemies was carried out along with collection of adult arthropods through sweep net method, and adopting the methods suggested by Reissig et al., [8] and Bayot et al., [9]. The relative abundance were calculated by Relative abundance of dominant species of pollinator was worked out by using the following formula given by Nath and Virakthamath [10] after pooling all the data.

Relative abundance of species A = (Number of particular species / Total no. of Insect species) X 100

2.1 Observation, Identification of Free Living Arthropods

Free-living insects were collected from Germination/seedling to siliqua maturation stages using a sweep net having a diameter of 30 cm, mesh size of 1.5 mm and attached with a 1.5 m long rod. Sweeping was done weekly in between 09.00 to 11.00 h of the day, and each sample was consisted of 30 sweeps encompassing an area from ground level to the top of the plants. The pollinators were collected during bloom of the crop. The collected insects were brought to the Entomology Laboratory of MRCFC Khudwani for identification and counting. "They were killed by storing in a freezer for a few hours, mounted on points, dried and morphotyped. Insects were identified up to species or genus level and also separated as pest, predator, pollinator and unknown categories" [11-15].

3. RESULTS AND DISCUSSION

Six insect- pests were recorded during the cropping season at different stages of *Brassica* species of which 3 belonged to Hemiptera, 1 to Lepidoptera, one each to the order Coleoptera and Diptera in kashmir valley (Table 1). It was observed that the insect- pest species infested different parts viz., leaves, shoot, flower and pod. Major pest which caused significant damage was *Cabbage butterfly*, *Leaf miner* and *Mustard aphid*. During the course of investigation, three natural enemies' viz., *Coccinella septempunctata*, *Apanteles glomeratus* and *Diaeretiella rapae* were found to be associated with insect-pest of rapeseed-mustard irrespective to the insect pest (Table 3). The abundance of insect pollinators recorded from mustard field during bloom of same experiment revealed 6

insect pollinators (Table 2) belonged to 3 orders. The highest mean relative abundance of *Apis cerana* (11.71%) followed by *Apis mellifera* (7.04%) and least 3.74% of *Erstalis arbustorum*. The Photographs of insect pests pollinators are listed in Table 2 and Fig. 1. Table 4 showed that the abundance, richness and diversity of pest, predator, pollinator and other category of insects varied from 0.8 ± 0.1 to 7.0 ± 1.6 , 0.3 ± 0.1 to 0.5 ± 0.2 and 0.5 ± 0.3 to 6.4 ± 1.6 per 30 sweeps, respectively and the results differed significantly. The richness and abundance of pollinators were significantly higher compared to pests, predators and other category. The pests revealed the highest diversity compared to predators, pollinators and other category. Several studies have shown that different insect pests infest rapeseed-mustard at different locations in India. Rai [16] reported "a total of 24 insect pests from India, out of which *L. erysimi*, *A. lugens proxima* and *B. Cruciferarum* were referred as the major pests while flea beetles, *P. Cruciferae* Goeze and cabbage butterfly, *P. Brassicae* L., *M. Persicae* Sulzer and *P. Xylostella* Lin. were considered as minor pest". "The total number of insect pests on rapeseed-mustard increased and stood at 38, of which 15 are considered economically important" (Bakhetia and Sekhon, 1984). "In 1991, the number of insect pests increased to 40" [17]. "Present, more than 43 species of insect pests have been reported to infest rapeseed mustard in India, in which about a dozen species are considered as major pests" [18]. Choudhury and Pal (2006) reported "fourteen insect including natural enemies on rapeseed-mustard. Out of which, *L. erysimi* was categorized as major pest". Singh et al. [19] reported "21 insect species during the various crop stages". "Nowadays, mustard aphid is observed as a national insect-pest" [20].




Fig. 1. Relative abundance of Insect Pollinators in rapeseed bloom

Table 1. Insect pest complex associated with mustard in MRCFC Khudwani Kashmir during rabi 2020-21

Common name	Scientific name	Order	Family	Feeding site	% Incidence		Mean	Status
					2020	2021		
Mustard aphid	<i>Lipaphis erysimi</i> (Kalt.)	Aphididae	Hemiptera	Leaf/Flowering stage	7.69	8.69	8.19	+++
Cabbage butterfly	<i>Peries brassicae</i>	Lepidoptera	Pieridae	Leaf	11.53	10.54	11.03	+++
Flea beetle	<i>Phyllotreta cruciferae</i> (Goeze)	Coleoptera	Chrysomelidae	Leaf	5.76	6.22	5.99	++
Pea leaf miner	<i>Phytomyza horticola</i>	Diptera	Agromyzidae	Leaf	9.61	10.69	10.15	++
Cabbage aphid	<i>Brevicoryne brassicae</i> (Linn.)	Hemiptera	Aphididae	Leaf/Flowering stage	7.22	8.12	7.67	+++
Shield Bug	<i>Carpocoris</i> sp.	Pentatomidae	Hemiptera	leaf	0.34	0.24	0.29	+
C.D.(P=0.05)					2.34	2.67		

Table 2. Insect Pollinator complex of mustard in MRCFC Khudwani Kashmir during rabi 2020-21

Scientific Name	Order	Family	Relative abundance (%)		Mean
			2020	2021	
<i>Apis mellifera</i>	Hymenoptera	Apidae	6.54	7.55	7.04
<i>Apis cerana</i>	Hymenoptera	Apidae	11.21	12.21	11.71
<i>Eristalis arbustorum</i>	Diptera	Syrphidae	3.73	3.76	3.74
<i>Lassioglossum mariginatum</i>	Hymenoptera	Halictidae	7.47	7.44	7.45
<i>Adrena</i> spp.	Hymenoptera	Andrenidae	6.54	6.04	6.29
<i>Eristalis tenax</i>	Diptera	Syrphidae	5.60	5.78	5.69
C.D.(P=0.05)			2.97	1.89	

Table 3. Natural enemies associated with insect-pests of rapeseed-mustard in MRCFC Khudwani during rabi 2020-21

Scientific name	Order and family	Host	Relative abundance	Economic status
<i>Coccinella septempunctata</i>	Coleoptera Coccinellidae	<i>Lipaphiserysimi</i> , <i>Myzuspersicae</i> , <i>Brevicorynebrassicae</i>	+++++	Major
<i>Apanteles glomeratus</i>	Hymenoptera Braconidae	<i>Pierisbrassicae</i>	++	Low
<i>Diaeretiella rapae</i>	Hymenoptera Braconidae	<i>Lipaphiserysimi</i> , <i>Myzuspersicae</i> , <i>Brevicorynebrassicae</i>	++	Low

Table 4. Average richness, abundance and diversity of arthropod fauna of mustard

	Pests	Natural Enemy (Parasitoids/predators)	Pollinators
Abundance	7.0 ± 1.6	1.0 ± 0.5	3.0 ± 8.5
Richness	1.8 ± 0.43	0.5 ± 0.2	6.4 ± 1.6
Diversity	0.8 ± 0.1	0.3 ± 0.1	0.5 ± 0.3



Fig. 2. Some insect pests of rape seed mustard In Kashmir Valley. 1, Shield bug, 2, *Lipaphis erysimi* 3, *Pieris brassicae* 4, *Phytomyza horticola*

4. CONCLUSION

In conclusion from the present investigation it can be concluded that the mustard aphid, flea beetle and cabbage butterfly were key pests of mustard found abundantly at different developing stages of the crop. Fortunately a number of natural enemies of these pests also present along with them. These beneficial bio control agents are useful for pest suppression in an ecologically viable and sustainable pest management programme.

ACKNOWLEDGEMENT

This work has been conducted under RCM Project sponsored by SKUAST-k at Mountain Research Centre for field crops Khudwani Kulgam.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Patel S, Singh C. Development of weather based forewarning model for mustard aphid, *Lipaphis erysimi* (Kalt.) in Tarai region of Uttarakhand. J. Agrometeorol. 2017a;19:382.
- Patel S, Singh C. Seasonal dynamics of *Lipaphis erysimi* (Kalt.): Arrival, peak and migration pattern in tarai region of Uttarakhand. J. Exp. Zool. India. 2017b; 20:1573-75.
- Patel S, Singh CP, Yadav SK. Seasonal incidence of mustard flea beetle, *Phyllotreta cruciferae* on *Brassica* species in relation to weather parameters at different dates of sowing. J. Ent. Zool. Stud. 2017a;5:673-77.
- Patel S, Yadav SK, Singh CP. The incidence of painted bug, *Bagrada hilaris* (Burmeister) on *Brassica* spp. and *Erucasativa* with respect to the date of sowing. J. Ent. Zool. Stud. 2017b;5:774-76.
- Patel S, Yadav SK, Singh CP. Bio-efficacy of insecticides against *Lipaphis erysimi* (Kalt.) in mustard ecosystem. J. Ent. Zool. Stud. 2017c;5:1247-50.
- Patel S, Yadav SK, Singh CP. Foraging behaviour of mustard sawfly, *Athalia lugens proxima* Klug. on rapeseed-mustard. J. Experim. Zool. 2018;21:379-81.
- Patel S, Yadav SK, Singh CP. Foraging behaviour of mustard sawfly, *Athalia lugens proxima* Klug. on rapeseed-mustard. J. Experim. Zool. 2018;21:379-81.
- Reissig WH, Heinrichs EA, Litsinger JK, Moody K, Fielder L, Mew TW, et al. Illustrated guide to Integrated Pest Management in Rice in Tropical Asia, International Rice Research Institute, Los Banos, The Philippines. 1986;411.
- Bayot RG, Cayabyab BF, Benigno EA, Ebuenga MD, Penalba FE. Monitoring populations of the brown plant hopper, *Nilaparvata lugens* (Stal), the green leafhopper, *Nephotettix virescens* (Distant) and spiders in rice at Maquiapo, Guagua, Pampanga. Philippines Entomologist. 1990;8(2):779-786.
- Nath S, Virathamath S. Pollinator fauna of sunflower and their relative abundance. Karnataka Journal of Agriculture Science. 2010;23(3): 517-518.
- Sharma HC, Rao MV. (Eds). National Seminar in Changing Scenario in Pest and Pest Management in India held during. 31 Jan - 1 Feb:Hyderabad, pp. 56-59.
- Yadav SK, Patel S. Insecticidal and repellent activity of some plant extracts against *Myzus persicae* (Sulzer) and *Brevicoryne brassicae* (Linnaeus). J. Ent. Zool. Stud. 2017;5:1434-39.
- Sollai G, Biolchini M, Crnjar R. Taste sensitivity and divergence in host plant acceptance between adult females and larvae of *Papiliohospiton*. Insect Sci. 2018;25:809–822.
- Feeny P, Städler E, Åhman I, Carter M. Effects of plant odor on oviposition by the black swallowtail butterfly, *Papilio polyxenes* (Lepidoptera: Papilionidae). J. Insect Behav. 1989;2:803–827.
- Patel S, Singh CP, Yadav SK. Monitoring of insect-pest complex on rapeseed-mustard at Pantnagar. Journal of Entomological Research. 2019;43(1):73-6.
- Rai BK. Pests of oilseed crops in India and their control. ICAR, New Delhi. 1976;121.
- Bakhetia DRC, Arora R. Changing scenario of insect pests and pest management in rapeseed mustard. In: Pests and Pest Management in India - The Changing Scenario; 1993.
- Purwar JP, Singh RK, Mall P. Eco-friendly management of insect pests in rapeseed-mustard. Indian Farmers Digest. 2004;37: 34-35.
- Singh R, Singh D, Rao VUM. Effect of abiotic factors on mustard aphid (*Lipaphis erysimi* Kalt.) on Indian Brassica. Indian J. Agric. Res. 2007;41:67-70.
- AICRP. Annual Progress Report. National Research Centre on Rapeseed-Mustard, Bharatpur - 321303, Rajasthan, India; 2015.

© 2023 Sofi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105091>