



# **Effects of Months of Coppicing on Cacao (*Theobroma cacao* L.) Chupons' Flowering and Yield in Edo State, Nigeria**

**Adejobi, K. B <sup>a\*</sup>, Adeosun, S. A <sup>a</sup>, Ibe, O. <sup>a</sup>,  
Ayegboyin, K. O. <sup>a</sup> and Famaye, A. O. <sup>a</sup>**

<sup>a</sup> *Agronomy and Soils Division, Cocoa Research Institute of Nigeria, P.M.B 5244, Ibadan, Nigeria.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ijpss/2025/v37i125877>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/148340>

**Original Research Article**

**Received: 20/09/2025**

**Accepted: 21/11/2025**

**Published: 11/12/2025**

## **ABSTRACT**

Rehabilitation through coppicing and chupon regeneration has been reported to improve cocoa yield. However, appropriate month for this operation is grossly scarce in literature. Therefore, investigation was carried out between 2017 and 2019 to evaluate effect of coppicing at different months on cocoa (*Theobroma cacao* L) chupons' flowering and yield performance in Cocoa Research Institute of Nigeria, Udonmora, Edo State. Months of coppicing considered as treatments were December 2017 to November 2018. The experiment was laid out in Randomized complete block design with three replications. Data were collected on number of flowers, number of cherelles, number of pods, fresh pod weight, fresh bean weight and dry bean weight of the chupons. Data were analyzed with ANOVA at  $P = .05$ . This study revealed that old cocoa trees

\*Corresponding author: E-mail: [jobikayode@gmail.com](mailto:jobikayode@gmail.com);

coppiced in the month of April were the most vigorous and showed earlier and steadier yield attributes than other treatments, coming into full flowering, cherelles and pods formation at 4, 9 and 14 months after coppicing, respectively. The old cocoa coppiced in April produced 8, 18, 27, 30 and 45 pods per tree (8999, 20189, 29631, 33331 and 50363 pods per hectare of coppiced cacao) after 14, 15, 16, 17 and 18 months, respectively. Also, it was observed that old cacao coppiced in the months of April and January produced the highest fresh pod weight, fresh bean weight and dry bean weight. April is therefore recommended as the appropriate month to coppice old cocoa plantation for yield performance of cocoa trees in Udonmora, Edo-State, Nigeria.

**Keywords:** *Chupon flowering; coppiced cocoa; old cocoa plantation; yield performance.*

## 1. INTRODUCTION

Cocoa is a tree crop which is among long list of commercially important species in tropical region of the world. It shares features of being perennial plant with long cycle and declining yields with age. It is also expensive to establish and has long waiting periods before starting to reap benefit of the harvest. Cocoa farmers are mostly driven into rehabilitation when there is fall in cocoa yield due to combined effects of plant ageing, reduction in cocoa population density linked to accumulated natural mortality and increased incidence of pest and disease (Aikpokpodion & Adeogun, 2011; Jagoret *et al.*, 2011; Mahrizal *et al.*, 2014; Akinnagbe, 2015; Wessel & Quist-Wessel, 2015; Adebisi *et al.*, 2018).

Coppicing is one of the perfected techniques of cocoa rehabilitation that CRIN (Cocoa Research Institute of Nigeria) has developed (Ofori *et al.*, 2022). It involves cutting down of the old cocoa tree in a slant position at 30 cm above ground level with a view to allow chupons growth, out of which the most vigorous and basal one or two are retained to replace the old cocoa tree. Somarriba *et al.* (2021) showed that regenerated amelonado trees in their sixth year after coppicing gave a yield of 1, 680 kg dry cocoa bean per hectare which is about four times the national average production of 460 kg per hectare. Other researchers got similar results on F3 Amazon where coppicing in November performed best in a monthly coppicing trial. Low bean yield of 400 kg/ha as obtainable in most cocoa plantations in Nigeria has been due to old age of the plantations and poor management practices. Adeyemi (1999) also got yield improvement under rehabilitated plot with soil application of NPK fertilizer. However, economic yield could only be evaluated after six years. In spite of these reported achievements on coppicing or chupon regeneration as a method of rehabilitation in cocoa, many cocoa farmers in

Nigeria are still reluctant in adopting the technology because it involves cutting down of their cocoa trees, accompanied with high cost implication and not wanting to miss a year of harvest on coppiced cocoa plantation (Osei-Bonsu *et al.*, 2002). For these reasons in Nigeria, Cocoa Rehabilitation Project (CRP) through coppicing and other rehabilitation techniques should be Federal Government initiative and part of the Economic Recovery Programme (ERP) package under the supervision of the World Bank (WB) and the International Monetary Fund (IMF) in conjunction with other institutions like Cocoa Research Institute of Nigeria (CRIN) and Agricultural Development Bank, as it is being done in Ghana and some other cocoa producing nations of the world (Kwaw-Nimesson & Tran, 2019). Although rehabilitation through coppicing and chupon regeneration has been reported to improve yield (Chibinga *et al.*, 2016; Adebisi *et al.*, 2018; Somarriba *et al.*, 2020; Adebayo *et al.*, 2022), but information on the appropriate month in a year to carry out this productive operation is grossly scarce in the literatures. Therefore, the objective of the study was to study the effect of month of coppicing on cocoa flowering and yield performance in Edo State, Nigeria.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

Field experiment was carried out at the experimental farm of Cocoa Research Institute of Nigeria (CRIN), Udonmora Station in Edo State between 2017 and 2019, covering two consecutive rainy seasons and two dry seasons. The location lies on latitude 6° 5' N and longitude 5° 50' E, a derived savanna zone of Nigeria. The rain fall is between 1000 – 1500 mm per annum. The maximum temperature ranges between 26 to 35 °C with an average of about 30 °C, while minimum temperature ranges from 15 to 25 °C

with an average of 20 °C. Relative humidity is high during the rainy season, ranges from 50 to 85% with an average of 75%. There are seasonal variations in the values of relative humidity, which varies from 65 to 89% during the rainy season and 46 – 70% during the dry season. The rainy season which runs from April to October is characterized by heavy rain, low ambient temperature and high humidity; while the dry season runs from November to March and is characterized by little or no rain, high ambient temperature and very low humidity.

## 2.2 Acquisition and Preparation of Experimental Materials

About 50 x 30 m land area of old cacao plantation of about fifty years was selected for the experiment. Plantain suckers used as shade crop for regenerated chupons were collected from experimental plots in the Station. Other materials used for the experiment were: Chain saw for cutting down of the cacao trunks, Long ruler for measurement of cacao height to be coppiced, Vernier Caliper for measuring the stem girth, Red paint for covering the cut surfaces against pest infestation and disease infection. The experimental plot measuring 50 by 30 m were mapped out and the experiment was laid out with an average of 144 stands of cocoa trees.

## 2.3 Treatments and Experimental Design

The field experiment comprised four trees randomly selected for each treatment, making forty-eight (48) and one hundred and forty-four (144) cacao stands per block and experimental site, respectively. Two stands were eventually tagged per treatment for data collection. The experiment was laid in Randomized Complete Block Design (RCBD) with three replications. The months of coppicing considered as treatments were: December, 2017 to November, 2018 to make up to twelve treatments based on the number of months in a calendar year. One hundred and forty-four (144) plantain suckers were planted at 3 x 3 m spacing as shade crop. The experiment was monitored for 22 months after coppicing. Measurement, pegging and holing for plantain suckers were carried out before coppicing. The coppicing was carried out on treatment monthly basis by complete removal of the main stems of cocoa trees to encourage the regeneration of the canopy by chupon growth. The 30 cm from the soil surface of cocoa trees to be coppiced was marked out with red

paint. Chain saw was placed on the marked portion of the trunk ready for coppicing in a slant position. Old cacao trunk was cut down in a slant position 30 cm above the soil surface. The cut surfaces were painted with red paint (Red paint contains red oxide) to prevent infestation and infection of pest and disease. A month after coppicing, there was chupon growth which was left for three (3) months before thinning. Three month after coppicing, the chupons on cocoa stump were pruned to two strong basal shoots. The two shoots were retained and allowed to develop mature canopies.

## 2.4 Data Collection

Data were collected on the yield parameters of cacao chupons. Number of flowers, number of cherelles and number of pods per chupon were determined by visual count; fresh pod weight, fresh bean weight and dry bean weight per chupon, were measured using weighing balance. These yield parameters were taken monthly for 22 months commencing from 4 months after coppicing (MAC).

Data collected were analysed using analysis of variance (ANOVA), and significant means separated by Duncan Multiple Range Test (DMRT) ( $P = .05$ ).

## 3. RESULTS AND DISCUSSION

Effects of months of coppicing on yield performance of cocoa are presented in Tables 1 - 6. Only old cocoa coppiced in the month of April produced flowers at 4 and 5 months after coppicing. Month of April gave significant ( $P = .05$ ) higher number of flowers at 4, 5, 6, 7 and 8 MAC (months after coppicing) compared to other months of coppicing. The significant higher number of flowers recorded in the month of April was distant followed by old cocoa coppiced in the months of June, July, March, May and September at 8 MAC (Table 1).

The results of Tables 2 and 3 also showed that coppicing in the month of April enhanced the number of cherelles and pods when compared to other treatments. It was the most vigorous, and showed early and steady yield attributes than other treatments, coming into full flowering, cherelles and pods formation as early as 4, 9 and 14 months after coppicing (Tables 1, 2 & 3). Therefore, the old cocoa coppiced in April was able to produce 8, 18, 27, 30 and 45 pods per tree after 14, 15, 16, 17 and 18 MAC, respectively.

**Table 1. Effects of months of coppicing on number of flowers per hectare at 4, 5, 6, 7 & 8 MAC**

| Treatments | Months after coppicing |         |         |         |          |
|------------|------------------------|---------|---------|---------|----------|
| Months     | 4                      | 5       | 6       | 7       | 8        |
| January    | 0.00b                  | 0.00b   | 0.00b   | 0.00c   | 0.00c    |
| February   | 0.00b                  | 0.00b   | 0.00b   | 0.00c   | 0.00c    |
| March      | 0.00b                  | 0.00b   | 23,320a | 10,360b | 10,360b  |
| April      | 10,350a                | 26,290a | 23,330a | 49,99a  | 102,570a |
| May        | 0.00b                  | 0.00b   | 0.00b   | 4,077b  | 7,770b   |
| June       | 0.00b                  | 0.00b   | 0.00b   | 0.00c   | 18,880b  |
| July       | 0.00b                  | 0.00b   | 0.00b   | 0.00c   | 11,111b  |
| August     | 0.00b                  | 0.00b   | 0.00b   | 0.00c   | 0.00b    |
| September  | 0.00b                  | 0.00b   | 0.00b   | 0.00b   | 6,290b   |
| October    | 0.00b                  | 0.00b   | 0.00b   | 0.00b   | 0.00c    |
| November   | 0.00b                  | 0.00b   | 0.00b   | 0.00b   | 0.00c    |
| December   | 0.00b                  | 0.00b   | 0.00b   | 0.00b   | 0.00c    |

Means followed by the same letters in each column are not significantly different by Duncan Multiple Range Test (DMRT) ( $P = .05$ )

**Table 2. Effects of months of coppicing on number of cherelles per hectare at 9, 10, 11, 12 & 13 MAC**

| Treatments | Months after coppicing |         |         |          |         |
|------------|------------------------|---------|---------|----------|---------|
| Months     | 9                      | 10      | 11      | 12       | 13      |
| January    | 10.00c                 | 16,630a | 18,221a | 22,554a  | 22,554a |
| February   | 4.00c                  | 8,855b  | 11,077b | 14,410bc | 13,521b |
| March      | 4,077cd                | 8,477b  | 9,588b  | 11,077c  | 12,555b |
| April      | 21,109a                | 16,254a | 20,332a | 21,409a  | 23,142a |
| May        | 1,111d                 | 14,399a | 18,110a | 20,332ab | 23,398a |
| June       | 6,666bc                | 9,222b  | 11,444b | 12,555c  | 15,521b |
| July       | 3,699cd                | 9,222b  | 11,810b | 12,555c  | 14,410b |
| August     | 10.00c                 | 8,477b  | 11,444b | 10,333c  | 12,555b |
| September  | 7.00c                  | 8,477b  | 11,077b | 10,333c  | 11,810b |
| October    | 8.00c                  | 8,477b  | 11,810b | 11,810c  | 11,188b |
| November   | 2.00c                  | 8,477b  | 11,444b | 11,810c  | 12,188b |
| December   | 11.00c                 | 8,855b  | 12,555b | 11,077c  | 12,921b |

Means followed by the same letters in each column are not significantly different by Duncan Multiple Range Test (DMRT) ( $P = .05$ )

**Table 3. Effects of months of coppicing on number of pods per hectare at 14, 15, 16, 17 & 18 MAC**

| Treatments | Months after coppicing |         |         |         |          |
|------------|------------------------|---------|---------|---------|----------|
| Months     | 14                     | 15      | 16      | 17      | 18       |
| January    | 4,810b                 | 9,254b  | 12,587b | 15,554b | 34,998b  |
| February   | 4,477b                 | 2,222c  | 3,333d  | 4,444cd | 19,810d  |
| March      | 3,333b                 | 5,921bc | 7,777c  | 7,410c  | 25,743c  |
| April      | 8,999a                 | 20,187a | 29,631a | 33,331a | 50,363a  |
| May        | 1,477b                 | 2,588c  | 2,588d  | 2,588d  | 19,444d  |
| June       | 2,222b                 | 2,966c  | 3,699d  | 3,333cd | 19,444d  |
| July       | 4,077b                 | 2,588c  | 4,077d  | 3,333cd | 19,444d  |
| August     | 2,222b                 | 2,966c  | 3,333d  | 4,444cd | 19,810d  |
| September  | 2,222b                 | 2,588c  | 3,333d  | 5,188cd | 19,188d  |
| October    | 2,222b                 | 3,333c  | 2,966d  | 5,188cd | 19,188d  |
| November   | 1,855b                 | 2,966c  | 3,699d  | 3,333cd | 19,555d  |
| December   | 2,588b                 | 2,966c  | 3,333d  | 4,077cd | 21,299cd |

Means followed by the same letters in each column are not significantly different by Duncan Multiple Range Test (DMRT) ( $P = .05$ )

These pod yields translated to 8,999, 20,189, 29,631, 33,331 and 50,363 pods per hectare of coppiced cocoa plantation, respectively (Table 4). Also, it was clearly observed that old cocoa coppiced in the months of April and January produced the highest fresh pod weight, fresh bean weight and dry bean weight (Tables 4, 5 & 6). It is worthy of note that at 18 MAC, month of July gave the lowest dry bean weight of 484 kg/ha while the highest dry bean weight of 4,654 kg/ha was obtained from the month of April (Table 6). This implies that the low bean yield of 400 kg/ha as obtainable in most cocoa plantation in Nigeria has been attributed to old age of the plantation and poor management practices by the farmers. Several researchers reported that the highest cocoa yield are achieved between 15

and 25 years, while Krung and Quartey- Papafio (1964) after conducting cocoa survey throughout the growing regions of the world recommended 30 – 40 years as the average economic life span of cocoa trees. The highest dry bean weight of 4,654 kg/ha obtained at 18 MAC for the month of April was far beyond 1,800 kg/ha dry bean of cocoa recorded after 18 month of coppicing. Also, Somarriba et al. (2021) noted that regenerated Amelonado trees in their sixth year after coppicing gave a yield of 1,680 kg/ha which was about four times the national average of 460 kg/ha at that time. The relative low yields recorded by the two authors could be as a result of the fact that appropriate techniques and period of application were not taken into consideration to get the required results. Adebisi et al. (2018)

**Table 4. Effects of months of coppicing on fresh pod weight (kg/ha) at 14, 15, 16, 17 & 18 MAC**

| Treatments | Months after coppicing |         |         |          |           |
|------------|------------------------|---------|---------|----------|-----------|
| Months     | 14                     | 15      | 16      | 17       | 18        |
| January    | 8,069a                 | 10,703a | 17,654a | 29,000a  | 45,987a   |
| February   | 1,938b                 | 1,776c  | 2,664c  | 2,840d   | 11,549c   |
| March      | 1,265ab                | 1,570bc | 3,040bc | 3,841bcd | 15,751abc |
| April      | 7,925a                 | 9,595a  | 16,501a | 29,610a  | 60,407a   |
| May        | 169.0b                 | 266.0c  | 722.0bc | 1,280bc  | 10,317bc  |
| June       | 1,029ab                | 1,894bc | 2,380bc | 9,468ab  | 10,356abc |
| July       | 932.0b                 | 840bc   | 1,324bc | 1,099cd  | 9,444bc   |
| August     | 788ab                  | 882.0bc | 762c    | 1,380cd  | 7,707bc   |
| September  | 992.0ab                | 805.0bc | 2,400bc | 2,130cd  | 10,002c   |
| October    | 666.0ab                | 2,571bc | 2,571bc | 4,968cd  | 10,862bc  |
| November   | 659.0ab                | 882.0bc | 4,886bc | 4,126cd  | 13,286abc |
| December   | 2,385ab                | 1,334bc | 1,487bc | 1,334cd  | 13,205abc |

Means followed by the same letters in each column are not significantly different by Duncan Multiple Range Test (DMRT) ( $P < 0.05$ )

**Table 5. Effects of months of coppicing on fresh bean weight (kg/ha) at 14, 15, 16, 17 & 18 MAC**

| Treatments | Months after coppicing |         |         |         |         |
|------------|------------------------|---------|---------|---------|---------|
| Months     | 14                     | 15      | 16      | 17      | 18      |
| January    | 2,688a                 | 3,576a  | 5,876a  | 9,908a  | 13,246a |
| February   | 532b                   | 592bc   | 788b    | 845b    | 1,058b  |
| March      | 429ab                  | 523bc   | 1,013b  | 1,280ab | 1,917ab |
| April      | 2,641a                 | 3,198a  | 5,500ab | 9,987a  | 13,654a |
| May        | 70.0b                  | 90.0c   | 280.0b  | 475.0ab | 456.0ab |
| June       | 364.0b                 | 654.0bc | 795b    | 3,167ab | 1,452ab |
| July       | 320b                   | 290bc   | 445b    | 369b    | 484.0b  |
| August     | 262.0b                 | 294.0bc | 257.0b  | 469b    | 572.0b  |
| September  | 334ab                  | 271bc   | 850b    | 723b    | 669b    |
| October    | 223b                   | 859abc  | 859b    | 1,656ab | 954.0b  |
| November   | 220b                   | 294bc   | 1,630ab | 1,375ab | 478.0ab |
| December   | 798ab                  | 447.0bc | 495.0b  | 447ab   | 3,401ab |

Means followed by the same letters in each column are not significantly different by Duncan Multiple Range Test (DMRT) ( $P < 0.05$ )

**Table 6. Effects of months of coppicing on dry bean weight (kg/ha) at 14, 15, 16, 17 & 18 MAC**

| Treatments | Months after coppicing |         |         |         |         |
|------------|------------------------|---------|---------|---------|---------|
| Months     | 14                     | 15      | 16      | 17      | 18      |
| January    | 899.0a                 | 1,190a  | 1,959a  | 3,577a  | 4,400a  |
| February   | 148c                   | 222d    | 333d    | 355c    | 481c    |
| March      | 147b                   | 176c    | 3340c   | 429     | 639.0b  |
| April      | 889a                   | 1,266a  | 1,834a  | 3,629a  | 4,654a  |
| May        | 30c                    | 35d     | 95d     | 158c    | 152c    |
| June       | 123b                   | 98c     | 270c    | 1,056b  | 543.0bc |
| July       | 108b                   | 97.0cd  | 150c    | 124.0bc | 162.0bc |
| August     | 88.0bc                 | 98.0cd  | 87.0cd  | 157.0b  | 195.0bc |
| September  | 122.9b                 | 91.0cd  | 289.0c  | 245.0b  | 223.0bc |
| October    | 75.0bc                 | 287.0bc | 289.0c  | 557.0b  | 318.0bc |
| November   | 73.0bc                 | 99.0cd  | 547.0bc | 459.0b  | 1,667bc |
| December   | 267.0b                 | 149.0c  | 165.0c  | 150.0b  | 1,133b  |

Means followed by the same letters in each column are not significantly different by Duncan Multiple Range Test (DMRT) ( $P < 0.05$ )

also reported that the appreciable increase in the yield of coppiced trees is an indication that the vibrant young trees from the coppiced old trees is increasing in size and height and is capable of carrying more cocoa pods. He equally observed 78.4 mean pod production at fifth year after coppicing. This work has rightly confirmed that rehabilitation through coppicing and chupon regeneration significantly enhanced the yield performance of cocoa as earlier reported (Chibinga *et al.*, 2016; Adebiyi *et al.*, 2018; Somarriba *et al.*, 2020), but information on the appropriate month in a year to carry out this productive operation is grossly scarce in the literature and that is the essence of this study.

#### 4. CONCLUSION AND RECOMMENDATION

This work has confirmed that rehabilitation through coppicing and chupon regeneration significantly enhanced the yield performance of cocoa. The appropriate techniques and period of application should be taken into consideration to get the required results. To obtain results close to the ones recorded in this work in farmers' farms, more training of farmers through farmers' field school to enable them rehabilitate their farms by themselves should be embarked upon by research scientists. Therefore, month of April is recommended for coppicing in Edo State for optimum yield. However, the recommendation is location specific. This calls for this research to be carried out in all cocoa ecologies in Nigeria.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of this manuscript.

#### ACKNOWLEDGEMENTS

We are very grateful to Executive Director and Management of Cocoa Research Institute of Nigeria for their logistics support for this research.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Adebayo, S. T., Oyawole, F. P., Sanusi, R. A., & Afolami, C. A. (2022). Technology adoption among cocoa farmers in Nigeria: What drives farmers' decisions?. *Forests, Trees and Livelihoods*, 31(1), 1-12. <https://doi.org/10.1080/14728028.2021.2011789>
- Adebiyi, S., Uwagboe, E. O., Agbongiarhuoyi, A. E., Famuyiwa, B. S., Abdulkarim, I. F., & Williams, A. O. (2018). Effects of coppicing on yield of cocoa pods in Cocoa Research Institute of Nigeria (CRIN) demonstration plot. *International Journal of Development Research*, 08(06), 20941 <http://www.journalijdr.com/effects-coppicing-yield-cocoa-pods-cocoa-research-institute-nigeria-crin-demonstration-plot>
- Adebiyi, S., Uwagboe, E. O., Agbongiarhuoyi, A. E., Famuyiwa, B. S., Abdulkarim, I. F., & Williams, A. O. (2018). Effects of coppicing on yield of cocoa pods in Cocoa Research Institute of Nigeria (CRIN) demonstration

- plot. International Journal of Development Research, 08(06), 20941  
<http://www.journalijdr.com/effects-coppicing-yield-cocoa-pods-cocoa-research-institute-nigeria-crin-demonstration-plot>
- Aikpokpodion, P. O., & Adeogun, S. O. (2011). A diagnostic study of constraints to achieving yield potentials of cocoa (*Theobroma cacao* L.) varieties and farm productivity in Nigeria. *Journal of Agricultural Science*.  
<https://doi.org/10.5539/jas.v3n4p68>
- Akinagbe, O. M. (2015). Evaluation of constraints to implementation and adoption of cocoa resuscitation programmes in Southwest Nigeria. *Applied Tropical Agriculture*, 20(1), 42–47.  
<https://journals.futa.edu.ng/ata/index.php/ata/issue/view/11>
- Chibinga, O. C., Nyangito, M. M., Musimba, N. R. K., Simbaya, J., & Daura, M. T. (2016). Effect of coppicing levels on the re-growth of *Julbernardia globiflora*. *Livestock Research for Rural Development*, 28(3), Article 42.  
<http://www.lrrd.org/lrrd28/3/chib28042.htm>
- Jagoret, P., Michel-Dounias, I., & Malezieux, E. (2011). Long-term dynamics of cocoa agroforestry: A case study in central Cameroon. *Agroforestry Systems*, 86(3), 493–504. <https://doi.org/10.1007/s10457-011-9513-9>
- Krug, C. A., & Quartey-Papafio, E. (1964). World cocoa survey. *FAO Agricultural Studies No.* 63.  
<https://agris.fao.org/search/en/providers/122621/records/6473966a68b4c299a3fb4cc7>
- Kwaw-Nimeson, E., & Tian, Z. (2019). Impact assessment of the cocoa rehabilitation project on cocoa exports in Ghana. *Open Journal of Social Sciences*, 7(9), 207–219.  
<https://doi.org/10.4236/jss.2019.79015>
- Mahrizal, L., Nalley, L. L., Dixon, B. L., & Popp, J. S. (2014). An optimal phased replanting approach for cocoa trees with application to Ghana. *Agricultural Economics*, 45(3), 291–302.  
<https://doi.org/10.1111/agec.12065>
- Ofori, A., Padi, F. K., Ameyaw, G. A., Dadzie, A. M., Opoku-Agyeman, M., Domfeh, O., & Ansah, F. O. (2022). Field evaluation of the impact of cocoa swollen shoot virus disease infection on yield traits of different cocoa (*Theobroma cacao* L.) clones in Ghana. *Plos one*, 17(1), e0262461. doi: 10.1371/journal.pone.0262461
- Osei-Bonsu, K., Opoku-Ameyaw, K., Amoah, F. M., & Oppong, F. K. (2002). Cacao-coconut intercropping in Ghana: agronomic and economic perspectives. *Agroforestry Systems*, 55(1), 1–8.  
<https://doi.org/10.1023/A:1020271608483>
- Somarriba, E., Peguero, F., Cerda, R., Orozco-Aguilar, L., López-Sampson, A., Leandro-Muñoz, M. E., ... & Sinclair, F. L. (2021). Rehabilitation and renovation of cocoa (*Theobroma cacao* L.) agroforestry systems. A review. *Agronomy for Sustainable Development*, 41(5), 64.  
<https://doi.org/10.1007/s13593-021-00717-9>
- Somarriba, E., Peguero, F., Cerda, R., Orozco-Aguilar, L., López-Sampson, A., Leandro-Muñoz, M. E., Jagoret, P., & Sinclair, F. L. (2021). Rehabilitation and renovation of cocoa (*Theobroma cacao* L.) agroforestry systems. A review. *Agronomy for Sustainable Development*  
<https://doi.org/10.1007/s13593-021-00717-9>
- Wessel, M., & Quist-Wessel, P. M. F. (2015). Cocoa production in West Africa, a review and analysis of recent developments. *NJAS-Wageningen Journal of Life Sciences*, 74–75, 1–7.  
<https://doi.org/10.1016/j.njas.2015.09.001>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
 The peer review history for this paper can be accessed here:  
<https://pr.sdiarticle5.com/review-history/148340>